



Estimation and Comparison of Growth Curve in Broilers Through the Artificial Neural Networks and Gompertz Models

Emine BERBEROĞLU^{1*} Nagihan ÖZKAN²

¹Tokat Gaziosmanpaşa University, Agricultural Faculty, Department of Animal Science, Tokat
(orcid.org/0000-0002-7318-2728)

²Tokat Directorate Of Provincial Agriculture And Forestry
(orcid.org/0000-0002-7748-3528)

* e-mail: emine.berberoglu@gop.edu.tr

Alındığı tarih (Received): 06.02.2020

Kabul tarihi (Accepted): 30.03.2020

Online Baskı tarihi (Printed Online): 09.07.2020

Yazılı baskı tarihi (Printed): 30.08.2020

Abstract: Mathematical models offer great convenience in estimating the variation in the growth of the living. The time-dependent change in weight and body sizes of the living can be estimated easily via mathematical models. In this study, the growth curves of Ross 308 broilers were compared through the Gompertz model, which explains growth best and the ANN model, which is assumed to be an alternative to this model. The model with high-estimated R^2 and low-estimated MSE, MAD, and MAPE values were considered as the best model. The criteria obtained from the ANN and Gompertz models are 5625 and 2950 for MSE, 0.27, and 0.17 for MAPE; 0.5 and 1.2 for MAD, respectively while R^2 values were observed as 0.99 in both models. MSE and MAPE values were observed lower compared to the Gompertz model.

Keywords: Artificial neural networks, broiler, Gompertz model, growth curve live weight.

Etlik Piliçlerde Büyüme Eğrisinin Yapay Sinir Ağları ve Gompertz Modeli ile Tahmin Edilmesi ve Karşılaştırılması

Öz: Canlının büyümesindeki değişimin tahminlenmesinde matematiksel modeller büyük kolaylıklar sağlamaktadır. Canlının ağırlık ve vücut ölçülerindeki zamana bağlı değişiminin tahmini matematiksel modeller ile kolay şekilde yapılabilir. Çalışmada Ross 308 etlik piliçlerinin büyüme eğrileri, büyümeyi en iyi açıklayan Gompertz modeli ile bu modele alternatif olabileceği düşünülen YSA modeli karşılaştırılmıştır. R^2 'sı yüksek; HKO, OMS ve OMYH değerleri düşük tahminlenen model en iyi model olarak dikkate alınmıştır. YSA ve Gompertz modelinden elde edilen kriterler sırasıyla HKO için 5625 ve 2950; OMYH için 0.27 ve 0.17; OMS değerleri ise 0.5 ve 1.2; R^2 değerleri ise her iki model için de 0.99 olarak gözlenmiştir. HKO ve OMYH değerleri Gompertz modeline göre daha düşük olarak gözlenmiştir.

Anahtar Kelimeler: Yapay sinir ağları, Gompertz modeli, etlik piliç, canlı ağırlık, büyüme eğrisi

1. Introduction

Growth refers to the increase in the weight and body size of the living over a certain time. Physiological growth begins with the formation of zygote after sperm fertilizes the egg (Çolak et al, 2006). Despite being a hereditary feature, growth can vary by race, species, and individuals, and these differences can be explained by mathematics (Emsen et al., 2004). In the interpretation of growth parameters, there may be differences according to the characteristics analyzed and the mathematical model used. Besides, in cases where the same characteristics are explained by different models, the obtained

parameters can better explain the same characteristics in one model, while not in another model (Köyceyiz, 2003).

In studies using nonlinear models, growth curve models serve a guide for further breeding research, and thus desired genetic and phenotypic characters can be achieved in selections to be done in breeding. Furthermore, the studies also reveal what sort of interaction can be possible between the obtained genotypes and environmental factors, which can be explained by growth curves (Yıldız et al., 2009).

The purposes of the use of growth curves in breeding can be summarized as follows; to show

the change of measures of certain properties over time, to decide whether there are differences in terms of the researched properties, to determine the effects of different forms of care and feeding on the development of animals, to estimate the variable that could not be measured in some periods, to be able to make forward-looking decisions based on the data obtained to determine the optimal slaughter age, to ground a basis for the selection studies on growth curves and growth rate, to determine the general health conditions of animals (Yakupoğlu, 1999; Doğan, 2003; Lambe et al., 2006; Yıldız et al., 2009; Çelikoğlu et al., 2014).

With the advancement in computer technology, interest in artificial neural networks (ANN) has increased in recent years. The use of ANN, which is widely used in many areas, has also increased in agriculture. ANN is also often preferred because this model produces easy solutions to complex and nonlinear problems (Gevrekçi et al., 2011).

ANN has been used in various studies conducted on animal production. ANN refers to computer software created by the imitation and development of the information processing system in the human brain to find solutions to various problems. Created through the sampling of an existing biological neural network in the human brain, ANN has the skills to memorize, learn, reveal the relationships between variables, produce new information about what they have learned and generalize. There are important relationships between statistical methods and ANN technology, which can be successfully applied in many fields (Yazıcı et al., 2007).

In this study, the growth curves of live weight in broilers were modeled with the Gompertz model and ANN, and these two models were compared.

2. Materials and Methods

Gaziosmanpaşa University Agricultural Application and Research Center for 0-6 weeks

of live weights obtained from 121 Ross 308 broiler chicks fed in individual cages data. The study was approved by the Animal Experiments Local Ethics Committee of Tokat Gaziosmanpaşa University (2018-HADYEK-23).

The Gompertz model used to model the growth curve, which is one of the methods in this study, was calculated with the following equation:

$$Y = Ae^{-\beta e^{-kt}} \quad (1)$$

where;

Y 6th week live weight in the equation, **A** asymptotic weight, **β** integration constant, **exp** natural logarithm base, **t** refers to the time, and **k** the rate of growth. Parameter A is the highest live weight that the animal can reach (Şahin et al., 2014). The initial values of β_1 and β_2 were calculated based on the following equations.

$$\beta_1 = \frac{(y_2 - y_1)/(t_1 - t_2)}{b_0} \quad (2)$$

$$\beta_2 = -\log_e y(0) + \log_e \beta_0 \quad (3)$$

where,

y_1 and y_2 are the highest and lowest values of the weight variable corresponding to the widest time interval of t_1 and t_2 ; b_0 is the initial value of the parameter β_0 (Narinç et al., 2009).

ANN used in this study is the networks formed by interconnecting neurons in various ways through the sampling of the way the biological nervous system works. These networks can store information, learn, and specify the relationship between data (Öztemel, 2003). ANN has general characteristics such as being nonlinear, learning, generalization, applicability, and fault tolerance. As ANN is modeling of biological neural networks, it is necessary to analyze the structure of the biological nervous system initially to understand its structure. Neurons, which are the building blocks of the biological nervous system, consist of four main parts. Figure 1 shows the these parts as nucleus, axon, dendrites, and synapses (Çayıroğlu, 2003).




Figure 1. Biological nerve cell structure
Şekil 1. Biyolojik sinir hücre yapısı

The smallest units, which constitute the basis of the working system of ANN, are called operational elements or artificial nerve cells. In the simplest form, as presented in Figure 2, the

artificial nerve cell structure consists of five main components, being inputs (weights X_i) (W_i), aggregate function (NET), activation function (F (NET)) and output.




Figure 2. Artificial nerve cell structure
Şekil 2. Yapay sinir hücre yapısı

The sigmoid function is generally used as an activation function that minimizes error in the multi-layer sensor model (Figure 1) which is widely used nowadays (Öztemel, 2003). The backpropagation algorithm also uses the sigmoid function that produces real values between 0 and 1 as the activation function. Due to the sigmoid function in the backpropagation algorithm, real outputs can be provided, even between two close values, thus the right decision can be made. Therefore, reverse learning is inevitable in solving nonlinear problems (Nabiyev, 2003).

$f(x) = \frac{1}{1+e^{-NET}}$, here refers to the NET input value of the element being under the NET process. This value is determined through the aggregate function (Öztemel, 2003).

As presented in Figure 4, ANN is the structures formed as a result of the interconnection of the artificial nerve cells and consists of three layers. These are the input layer, interlayer, and the output layer (Öztemel, 2003).




Figure 3. Sigmoid activation function model
Şekil 3. Sigmoid aktivasyon fonksiyon modeli




Figure 4. Artificial neural network structure
Şekil 4. Yapay sinir ağı yapısı

A large number of connections can be established between cells in separate layers while no connection is established between cells in the same layer (Yılmaz, 2015). The data from the outside is taken by nerve cells placed in the input layer and transmitted to the input layer, interlayer, and output layer, respectively (Yılmaz, 2015). The network itself is responsible for the operation that occurs in the interlayer. This layer is, in a sense, the intelligence part of the network, and a large number of neurons in that layer leads the network to memorization. The network is expected to learn rather than memorizing. In this way, the network will be able to tolerate any changes made through the learning process, otherwise, it will not be able to solve a knowledge-based problem in a small change (Nabiyev, 2003).

Learning systems in ANN are of three types; supervised, unsupervised, and reinforcement learning systems (Takma et al., 2012). In a supervised, learning system, output values corresponding to the input values given to the network are also given to the network. In the unsupervised learning system, only input data is displayed to the network while the expected output data is not displayed. In a reinforcement learning system, information is given to the network according to the results obtained at the end of each iteration (Çayıroğlu, 2003; Öztemel, 2003).

In this study, the feed-forward backpropagation supervised algorithm was selected as a network type. The reason for preferring the algorithm is that during the training

of the network when a forward connection is first established, the network obtained outputs corresponding to the input values by using its weights, and reverse propagation algorithm ensures backward rearrangement of the weights to mitigate error that occurs in the output layer. Furthermore, the Levenberg-Marquardt algorithm was preferred as the training function of the network since it significantly increases the learning speed. According to Kaastra et al (1996); networks with an interlayer are generally found to be successful in solving ANN problems (Aksu et al., 2016). The number of interlayers used in the network was taken as one because the over-preferred number of interlayers increases the complexity of calculation and extends the time. The number of neurons in the interlayer was determined by the method of trial and error (Takma, 2012; Yılmaz, 2015). The percentage (%) values used to determine the number of sampling during the training of the network were taken as 5 -10 -15 -20 -30 -35%, respectively within the limits of the program. Training, validation, and test set percentages were tried as 50 -60 -70 -80 -90%, respectively in order to determine the most accurate network model. Afterward, the network was trained. The network updated its weight to determine the desired output in response to the given inputs so that the error between the outputs of the network and the output observed by the network was calculated by the network, and the new weights of the network were regulated by this margin of error.

In practice, 21 different percentage distributions were used within the limitations of the program. Table 1 presents these percentage distributions and sampling numbers corresponding to percentages. The number of neurons in the interlayer was determined in 5 different ways, being the number of neurons (10), minus two (9-8) and plus two (11-12) which is automatically assigned by the program. In order to achieve the most accurate result, a total of 105 network models including 5 different neurons (21x5) were tested in 21 applications. As a result of the analysis, the model with the highest R^2 value in all sets, including training, testing, and

validation set, was chosen as the best model (Şahin et al., 2014). In addition, the MSE and R² values in the training, validation, and test sets were examined to determine whether the network completed the training by learning or

memorizing. - The network calculated low MSE in the training set and high MSE in test set; high R² in the training set and low R² in the test set, however, the network model only memorized (Aka et al., 2018).

Table 1. ANN data set distribution

Çizelge 1. YSA veri seti dağılımı

Training set (%)	Validation set (%)	Test set (%)	Training set (%)	Validation set (%)	Test set (%)	Training set (%)	Validation set (%)	Test set (%)
50	15	35	60	15	25	70	15	15
50	20	30	60	20	20	70	20	10
50	25	25	60	25	15	70	25	5
50	30	20	60	30	10	80	5	15
50	35	15	60	35	5	80	10	10
60	5	35	70	5	25	80	15	5
60	10	30	70	10	20	90	5	5

In this study, SPSS 17.0 for Windows was used for parameter estimation of the Gompertz model, and MATLAB statistical package programs for model estimation with ANN. Comparing the two models, the coefficient of determination (R²), mean absolute percentage error (MAPE), mean absolute deviation (MAD), and mean squared error (MSE) criteria were used. R² values were used to determine the best estimating ANN model. (Asilkan et al., 2009; Yavuz et al., 2013). The equations of these criteria are presented below.

$$R^2 = \frac{\sum(\hat{y}_t - \bar{y})^2}{\sum(y_t - \bar{y})^2} \quad (4)$$

$$MAD = \frac{\sum_{t=1}^n |y_t - \hat{y}_t|}{n} \quad (5)$$

$$MAPE = \frac{\sum_{t=1}^n \left| \frac{y_t - \hat{y}_t}{y_t} \right|}{n} * 100 \quad (y_t \neq 0) \quad (6)$$

$$MSE = \frac{\sum_{t=1}^n (\hat{y}_t - y_t)^2}{n} \quad (7)$$

In the equations, y_t is the value observed in t period, \hat{y}_t is the value estimated through the model for t period, n is the total number of observations.

MAPE statistics are recognized to be superior to similar methods as it is expressed as a percentage (Çuhadar et al., 2009, Yavuz et al., 2013). According to the MAPE categorization made by Witt and Witt (1992) and Levis (1982); models detected below 10% are considered “very good”, those between 10-20% as “good”, those

between 20 -50%. as “acceptable” and those above 50% as “faulty” (Çuhadar et al., 2009).

3. Results and Discussion

In ANN analysis based on MSE, R² and scatter plots among 105 models, the model with 12 neurons, 80% training, 15% validation, and 5% test set was detected as the best model. Table 2 presents MSE and R² values for this model and Figure 5 presents the scatter plot of the model. The R² values obtained from the model were found as 0.97, 0.97, 0.99 for training, validation and test sets, respectively. These R² values are the highest R² values among other models. Therefore, this model was selected as the best among all the models. 0.99 R² value detected for the test set also indicates that the model is a very good one. Considering the MSE value of the sets, it can be observed that the MSE in the training set was low and the R² value was high, which indicates that the selected model did not memorize but successfully completed the training.

Table 2. MSE and R² values of the best model determined by ANN analysis

Çizelge 2. YSA analizi ile belirlenen en iyi modele ait HKO ve R² değerleri

Data Set	(%)	MSE	R ²
Training Set	80	7 005	0.97
Validation Set	15	11 457	0.97
Test Set	5	5 625	0.99

The distribution of two variables together is examined through scatter plots. According to the distribution of points on the scatter plot, it can be determined whether there is a weak, strong, and complete relationship between variables (Alpar,

2013). The best result in the plot showing the overlap of the network's output and target values is possible with the maximum stack on the 45-degree curve (Kubat, 2015).




Figure 5. Regression scatter plots of the twentieth application.
Şekil 5. Yirminci uygulamaya ait regresyon saçılım grafikleri

The regression line drawn in the training and validation sets in the plots of the model determined by ANN analysis in Figure 5 coincides with the 45-degree line and the test set is very close to this value. Other applications do not have such a close overlap. No other network models were preferred along with scatter plots and R^2 values since some networks only memorize.

The estimated R^2 value for the ANN reported by Ahmadi et al., (2007) and Roush et al., (2006) is 0.99. These values reported in the literature are in line with the estimated value of the model.

Table 3. Parameters of the Gompertz model
Çizelge 3. Gompertz modeline ait parametreleri

Model	A	$S_{\bar{x}a}$	β	$S_{\bar{x}b}$	k	$S_{\bar{x}k}$
Gompertz	3999	585.7	4.6	0.39	0.05	0.008

The reason why estimated values were found different from other studies was the fact that the

The parameters of the Gompertz model used in the study are given in Table 3. The parameters A, β , and k were found as 3999, 4.6, and 0.05, respectively. Parameter A was found lower than the studies of Nariç et al. (2007) and Eleroğlu et al. (2016) and higher than Mohammed (2015). Parameter β was found similar to the studies of Şekeroğlu et al. (2013), Eleroğlu et al. (2016) and Mohammed (2015); lower than Nariç et al. (2007) Parameter k was found in line with the studies of Mohammed (2015) and Şekeroğlu et al. (2013); and lower than Nariç et al. (2007) and Eleroğlu et al. (2016).

live weight values were taken at different periods, maintenance and feeding conditions and trial periods were different. Live weight increase continues over time and changes the initial value of parameter A.

Growth curve graphs for the weekly live weight values estimated and observed from the specified models are presented in Figures 6 and

7. The growth curve estimated by the Gompertz model shows that the shape of the growth curves coincided with the ANN through which a deviation between the 21-42 days was estimated.




Figure 6. Change of observed and estimated values through the Gompertz model

Şekil 6. Gompertz modeli ile tahminlenen ve gözlenen değerlerin değişimi




Figure 7. Change of observed and estimated values through the ANN model

Şekil 7. YSA modeli ile tahminlenen ve gözlenen değerlerin değişimi

Comparison criteria for Gompertz and ANN model are presented in Table 4. In Table 4, MSE, MAPE, MAD, and R^2 were 2950, 1.2, 0.2, and 0.99 for the Gompertz model, respectively, and 5625, 0.5, 0.27, and 0.99 for ANN. In both models, R^2 was found equal, while the MAD was

high in Gompertz while the other criteria were low. According to the results of the study, despite the high difference in MSE, both models made very good estimations considering R^2 , MAD, and MAPE values according to the classification of Witt and Witt (1992) on MAPE values.

Table 4. Estimation criteria of Gompertz and ANN models**Çizelge 4.** Gompertz ve YSA modele ait tahmin kriterleri

Model	MSE	MAD	MAPE (%)	R ²
Gompertz	2950	1.2	0.17	0.99
ANN	5625	0.5	0.274	0.98

Comparing the estimations with other studies, MSE was found higher than the studies of Eleroğlu et al. (2016) Mohammed (2015), higher than the first stocking density of Şekeroğlu et al. (2013) and similar with the other studies. R² values reported by Şekeroğlu et al. (2013), Adenaike et al. (2017), Roush et al. (2006), Mohammed (2015), Eleroğlu et al. (2016) and Topal et al. (2008) coincide with the value of the R value in the present study. The high R² value indicates a very good estimation. MAPE and MAD values were lower than Roush et al. (2006). MAPE and MAD values of ANN were found lower than Roush et al. (2006) and Ahmadi et al. (2007).

4. Conclusion

In the study, it was observed that ANN and Gompertz models gave close results. R² value is the same in both models; MAPE and MAD values were close to MSE, whereas Gompertz model was lower. However, ANN estimated the live weight at the end of fattening, which is important in stockbreeding, with small errors than the Gompertz model.

The ANN and Gompertz models gave close results. In Gompertz model, individual parameters are estimated and averaged for each animal. This causes loss of time and raw data loss as the parameters are averaged. ANN is a very easy method. A single model is obtained from the data of all animals. It prevents loss of time and data. In addition, since ANN is not affected by the changes in the data set, it can be retrained even when new information occurs or when changes occur.

As a result, ANN can be used as an alternative to classic growth curve models since it yields close results to the Gompertz model, prevents data loss, adapts to changes, and can be retrained.

References

- Adenaike AS, Akpan U, Udoh JE, Wheto M, Durosaro SO, Sanda AJ and Ikeobi CON (2017). Comparative evaluation of growth functions in three broiler strains of nigerian chickens. *Tropical Agricultural Science*. 40(4): 611–620.
- Ahmadi H, Mottaghtalab M and Zadeh NN (2007). Group method of data handling- typeneural network prediction of broiler performance based on dietary metabolizable energy, methionine and lysine. *The Journal of Applied Poultry Research* 16(4): 494-501.
- Aka S ve Akyüz (2018). Yapay sinir ağları analizi ile süreçlerinin iyileştirilmesi. *Ege Akademik Bakış Dergisi*, 18(2): 261-271.
- Aksu N ve Uçan K (2016). Zaman ve konum girdileri kullanılarak yapay sinir ağlarıyla referans evapotranspirasyonun tahmin edilmesi. *El-Cezeri Fen ve Mühendislik Dergisi*, 3(2): 204-221.
- Alpar R (2013). Çok Değişkenli İstatistiksel Yöntemler. Detay Yayıncılık. Bizim Büro ve Basım Evi. Ankara. 886.
- Asilkan Ö ve Irmak S (2009). İkinci el otomobillerin gelecekteki fiyatlarının yapay sinir ağları ile tahmin edilmesi. *Süleyman Demirel Üniversitesi İktisadi ve İdari Bilimler Fakültesi Dergisi*.14(2): 375-391.
- Çayroğlu İ (2003). İleri Algoritma Analizi Ders Notu. Karabük Üniversitesi Mühendislik Fakültesi Karabük.
- Çelikoğlu K ve Tekerli M (2014). Pırlak kuzularında büyüme eğrilerini etkileyen genetik ve çevresel faktörlerin belirlenmesi ve eğri parametreleri yönünden baba koçların değerlendirilmesi. I. Bazı çevresel faktörlerin canlı ağırlığa ilişkin büyüme eğrilerine etkileri. *Afyon Kocatepe Üniversitesi Veterinerlik Fakültesi, Zootekni Anabilim Dalı, Lalahan Hayvan Araştırma Enstitüsü Dergisi*, 54(1): 8-14.
- Çolak C, Orman MN ve Ertuğrul O (2006). Simental x Güney Anadolu Kırmızısı sığına ait beden ölçüleri için basit doğrusal ve lojistik büyüme modeli. *Ankara Üniversitesi Veterinerlik Fakültesi Dergisi*, 53: 195-199.
- Çuhadar M, Güngör İ ve Göksu A (2009). Turizm talebinin yapay sinir ağları ile tahmini ve zaman serisi yöntemleri ile karşılaştırmalı analizi: Antalya iline yönelik bir uygulama. *Süleyman Demirel Üniversitesi İktisadi ve İdari Bilimler Fakültesi Dergisi*, 14(1): 99-114.
- Doğan İ (2003). Kuzularda büyümenin çok boyutlu ölçekleme yöntemi ile değerlendirilmesi. *Uludağ University Journal of The Faculty of Veterinary Medicine* 22(1-3): 33-37.
- Eleroğlu, H, Bircan, H, Yıldırım, A. ve Kılıç, F, (2016). Ticari etlik piliçlerde büyüme eğrilerinin doğrusal olmayan modeller kullanılarak karşılaştırılması.

- Tavukçuluk Araştırma Dergisi 13(2): 12-16.
- Gevrekçi Y, Yeğenoğlu ED, Akbaş Y ve Sesli M (2011). Yapay sinir ağlarının tarımsal alanda kullanımı. Ege Üniversitesi Ziraat Fakültesi Dergisi, 48(1): 71-76.
- Kaastra, I and Boyd M (1996). Designing a neural network for forecasting financial and econometric time series, *Neurocomputing*, 10: 215-236.
- Köyceyiz, F (2003). İvesi ve Morkaraman Kuzularında Değişik Vücut Ölçüleri Bakımından Büyüme Eğrileri., Atatürk Üniversitesi Fen Bilimleri Enstitüsü Yayınlanmamış Yüksek Lisans Tezi, Erzurum.
- Kubat, C (2015). Matlab Yapay Zeka ve Mühendislik Uygulamaları. *Deniz Ofset Matbaacılık*, s. 762, İstanbul.
- Lambe NR, Navajas EA, Simm G and Bunger L (2006). A Genetics investigation of various growth models to describe growth of lambs of to contrasting breeds. *Journal of Animal Science* 84: 2642-2654.
- Levis CD (1982). *Industrial and Business Forecasting Methods*, Butterworths Publishing. London.
- Mohammed FA (2015). Comparison of three nonlinear functions for describing chicken growth curves. *Science Agriculture*, 9(3): 120-123.
- Nabiye, VV (2003). Yapay Zeka. Birinci Baskı, Ankara. Seçkin Yayıncılık. 710.
- Nariç D, Aksoy T, Çürek D. ve Karama, E (2007). Farklı gelişme hızına sahip etlik piliçlerde büyümenin analizi. *Hayvancılık Araştırma Dergisi*, 17(2):1-8.
- Nariç D, Aksoy T, Karaman E ve Karabağ K (2009). Japon bıldırcınlarında yüksek canlı ağırlık yönünde uygulanan seleksiyonun büyüme parametreleri üzerine etkisi. *Akdeniz Üniversitesi Ziraat Fakültesi Dergisi*, 22(2): 149-156.
- Öztemel E (2003). Yapay Sinir Ağları. Birinci Baskı, İstanbul. *Papatya Yayıncılık*, 231.
- Roush WB, Dozier WA and Branton SL (2006). Comparison of gompertzandneural network models of broiler growth. *Poultry Science* 85(4): 794-797.
- Şahin A, Ulutaş Z, Karadavut U, Yıldırım A ve Arslan S (2014). Anadolu mandası malaklarında büyüme eğrisinin çeşitli doğrusal olmayan modeller kullanılarak karşılaştırılması. *Kafkas Üniversitesi Veterinerlik Fakültesi Dergisi*, 20(3): 357-362.
- Şekeroğlu A, Tahtalı Y, Sarıca M, Gülay MŞ, Abacı S. ve Duman M (2013). Farklı yerleşim sıklıklarındaki etlik piliçlerin büyüme eğrilerinin gompertz modeli ile karşılaştırılması. *Kafkas Üniversitesi Veterinerlik Fakültesi Dergisi*, 19(4): 669-672.
- Takma Ç, Atı, H ve Aksakal V (2012). Çoklu doğrusal regresyon ve yapay sinir ağı modellerinin laktasyon süt verimlerine uyum yeteneklerinin karşılaştırılması. *Kafkas Üniversitesi Veterinerlik Fakültesi Dergisi*, 18(6): 941-944.
- Witt SF and Witt C (1992). *Modeling and Forecasting Demand in Tourism*. Academic Pres, London. 195.
- Yakupoğlu Ç (1999). Etlik piliçlerde büyüme eğrilerinin karşılaştırılması. Ege Üniversitesi Fen Bilimleri Enstitüsü, Yayınlanmamış Yüksek Lisans Tezi, İzmir.
- Yazıcı AC, Ögüş E, Ankaralı S, Canan S, Ankaralı H ve Akkuş, Z (2007). Yapay sinir ağlarına genel bakış. *Türkiye Klinikleri Dergisi*, 27: 65-71.
- Yavuz, S ve Deveci M (2013). İstatiksel normalizasyon tekniklerinin yapay sinir ağı performansına etkisi. *Erciyes Üniversitesi İktisadi ve İdari Bilimler Fakültesi Dergisi*, 40: 167-187.
- Yıldız G, Soysal MI ve Gürcan EK (2009). Tekirdağ ilinde yetiştirilen Karacabey merinosu kıvrıcık melezi kuzularda büyüme eğrilerinin farklı modellerle belirlenmesi. *Tekirdağ Ziraat Fakültesi Dergisi*, 6(1): 11-19.
- Yılmaz B (2015). Akarçay Havzasında Çözünmüş Oksijen Değerlerinin Yapay Sinir Ağları ile Belirlenmesi. (Uzmanlık Tezi), Orman ve Su İşleri Bakanlığı Su Yönetimi Genel Müdürlüğü, Ankara