



Beekeeping Structure, Problems and Colony Losses in the Aegean Region of Turkey

Aytül UÇAK KOÇ^{1*} Mete KARACAOĞLU²

¹ Adnan Menderes University, Kocarli Vocational College, 09100 Aydın-Turkey

² Adnan Menderes University, Faculty of Agriculture, Animal Science, 09100 Aydın- Turkey

*e-mail: aucak@adu.edu.tr

Alındığı tarih (Received): 07.09.2016

Kabul tarihi (Accepted): 14.12.2016

Online Baskı tarihi (Printed Online): 22.12.2016

Yazılı baskı tarihi (Printed): 30.12.2016

Abstract: We aimed to determine the socioeconomic structure, colony management practices, colony losses and problems of beekeeping in the Aegean region of Turkey. According to the stratified sampling method, enterprises were divided into five groups according to the number of colonies. We determined that the average age of beekeepers was between 47.1±2.35 years (Group III) and 56.1±3.05 years (Group I). Beekeeping experience ranged from 14.4 ±24.2 years (Group I) and 26.4±2.91 years (Group V). The wintering losses of the groups between 2009 and 2012 varied from 6.0% (Group IV) to 29.4% (Group I) and the honey yield ranged from 11.4±2.30 kg (Group I) to 21.4±1.38 kg (Group V). The primary problems of beekeepers in the region are overcrowding of migratory beekeepers at certain times of the year, varroa mite and diseases and lower honey prices. In conclusion, a lower education level, an increased beekeeper age, concern about colony management, wintering with weak population and increasing varroa mite and diseases problems due mainly to overcrowding of migratory beekeepers in the region are some of the characteristics of beekeeping in the Aegean Region. These problems can be solved by reorganizing the baiting of beekeepers, providing support only to beekeepers that make a living from beekeeping (Group IV and V) and creating an organizational structure among beekeepers.

Keywords: Beekeeping, colony losses, honey yield, problems, questionnaire

Ege Bölgesi'nde Arıcılığın Yapısı, Koloni Kayıpları ve Sorunları

Öz: Bu çalışmada, Ege Bölgesinin sosyo-ekonomik yapısı, koloni yönetimindeki uygulamalar, koloni kayıpları ve arıcılık sorunlarının belirlenmesi amaçlanmıştır. Tabakalı örnekleme yöntemine göre, işletmeler koloni sayılarına göre beş gruba ayrılmıştır. Ankete katılan arıcıların ortalama yaşı, 47.1±2.35 (III. Grup) ile 56.1±3.05 (I. Grup) arasındadır. Ortalama arıcılık deneyimleri 14.4±2.24 (I. Grup) ile 26.4±2.91 yıl (V. Grup) arasında değişmiştir. Grupların 2009-2012 yıllarındaki ortalama kış kayıpları %6.0 (IV. Grup) ile %29.4 (I. Grup) arasında, bal verimi ortalaması 11.4±2.30 (I. Grup) ile 21.4±1.38 kg (V. Grup) olarak belirlenmiştir. Bölge arıcılarının başlıca sorunları, göçer arıcıların belli zamanda aşırı yoğunlaşması, varroa zararlısı ve balın düşük fiyata satılması olarak gösterilmiştir. Sonuç olarak, arıcıların eğitim seviyelerinin düşük, yaş ortalamalarının yüksek olduğu, koloni bakımına önem verdikleri, zayıf kolonilerle kışa girdikleri ve yöreye göçer arıcıların aşırı yığılmasından dolayı varroa ve bazı hastalıkların artış gösterdiği belirlenmiştir. Arıcıların sorunları, konaklama ile ilgili yeni düzenlemelerin yapılması, desteklemelerin sadece geçimini arıcılıktan sağlayanlara (IV. ve V. grup) verilmesi ve arıcılığın örgütlü bir yapıya kavuşması ile çözüme ulaşacaktır.

Anahtar Kelimeler: Arıcılık, koloni kayıpları, bal verimi, sorunlar, anket

1. Introduction

Beekeeping contributes to crop production via pollination and is one of the most commonly performed agricultural activities in the world, valued its honey, pollen, royal jelly, bee venom, propolis, beeswax, queens and production of

packaged honeybees. According to FAO statistics from 2014, China is the world's beekeeping leader with 9 million colonies; Turkey is in second with 7 million colonies. China is also the highest honey-producing country in the world with its production of 473600 tons. Turkey and

the United States follow with productions of 103525 and 80862 tons, respectively (Anonymous, 2016).

Because of the Mediterranean climate and the thick pine forests along the coastal regions, many Turkish beekeepers prefer living in the Aegean region of Turkey to benefit from its rich nectar and pollen sources (Karacaoğlu and Uçak Koç, 2007; Uçak Koç and Karacaoğlu, 2010). When the nectar and pollen sources are exhausted in May in the region, the beekeepers start migrating to the central regions and then to the northern and eastern regions of Turkey during the summer. In September, beekeepers move back to the *calabrian pine* (*pinus brutia*) forests because of the rich pine cochineal (*Marchalina hellenica*) in these forests in the Muğla, Aydın and İzmir provinces in the region. In autumn, nearly half of the Turkish beekeepers migrate to the region to winter their colonies there.

The ecological characteristics of the region resulted in the Muğla ecotype of the Anatolian bee (*Apis mellifera anatoliaca*). The Muğla ecotype is also known as the Aegean ecotype. In some studies, it has been reported that the morphological and physiological characteristics of the Aegean ecotype differ from those of other native honeybee genotypes in Turkey (Doğaroğlu et al., 1992; Güler and Kaftanoğlu, 1999; Gençer and Karacaoğlu, 2003; Karacaoğlu, 2005)

With the goal of attracting many beekeepers to migrate to the south of the Aegean region, it is beneficial to determine the problems, practices in colony management and wintering losses in the region. After the high colony losses seen in recent years in many countries, regular survey studies collecting information about colony losses, current beekeeping problems and drug usage information have become important (Danka and Villa, 2000; Caron and Hubner, 2001; Lawal and Banjo, 2010).

Aegean region of Turkey, with its climatic characteristics and rich flower and pine honey sources, is an important locale for migratory beekeepers and merits studies of the socio-economic characteristics, colony management and colony losses in recent years.

2. Materials and Methods

We relied on face-to face meetings with the beekeepers in Aydın province, Turkey. In September and November 2012, we visited 79 beekeepers and collected data via surveys. The questionnaire was composed of 115 questions related to personal, colony management, wintering losses, diseases and varroa mites, marketing and organization. In order to sample, 79 out of the 878 members of the Aydın Beekeepers Association (9%) were selected according to stratified random sampling. The beekeeping enterprises were divided into five groups according to the colony numbers: 9 beekeepers in group I (<50 colonies), 22 beekeepers in group II (51–125 colonies), 18 beekeepers in group III (126–200 colonies), 16 beekeepers in group IV (201–350 colonies) and 14 beekeepers in group V (>351 colonies). Chi-square test (Fisher's Exact Test) was used to analyze the data statistically.

3. Results and Discussion

The first group of beekeepers stated that they generally do beekeeping as a hobby and an interest for meeting the needs of their family and relatives. On the other hand, beekeepers in groups II and III do beekeeping for extra work to increase the income of their family. Approximately 80–85% of beekeepers in groups IV and V stated that they do beekeeping as a profession and added that their main income was beekeeping. About 70% of the beekeepers specified that the reason they started their beekeeping business was because of their father's occupation.

The average age, experience and number of family members of the beekeepers were determined to be 50.4 ± 1.20 years, 18.3 ± 1.09 years and 4.0 ± 0.14 persons, respectively.

In this study, even though the average ages of beekeepers in the different groups were similar, the number in the groups decreased as the colony number increased, but the experience of the beekeepers increased. Beekeepers in groups IV and V (colony numbers between 200 and 500) acquired beekeeping as a profession, producing flower and pine honey and pollen; these

individuals also want to improve their knowledge about the varroa mite and diseases.

The beekeepers generally record the age of the queen and renew the queens in the spring (April–May). In this study, even though changing the queen every two years is accepted as a good practice, the higher migration rate seen in groups IV and V results in aging queen bees quickly because of their higher rates of egg-laying almost over the course of the entire year. Due to the higher migration rate seen in groups IV and V, changing the queen honeybees every year may be advised.

The local genotype of the region, the Aegean ecotype of the Anatolian bee, predominated among the beekeepers. In addition, the preferences for Italian and Caucasian genotypes increased depending on the increase in colony numbers (13% in group II, 33% in group III, 44% in group IV and 79% in group V).

The honey yield and sugar consumption averages of the groups are listed in Table 1. We

determined that the differences among the group averages in terms of honey yield were statistically significant ($P < 0.01$). Group I was similar to group II, but differed from groups III, IV and V ($P < 0.05$). Group II was also different from groups IV and V ($P < 0.05$). The average sugar consumption among the groups, on the other hand, was found to be statistically similar ($P > 0.05$) and the average sugar consumption per colony changed from 6.5 kg to 7.9 kg. The beekeepers in groups I and II usually stopped over within the boundaries of Aydın and Muğla provinces over the entire year, but groups III, IV and V typically changed their locations 4–5 times per year. Beekeepers in groups III, IV and V moved to the north of the Aegean region and the central Anatolia region; sometimes, they went to Thrace region to benefit from the sunflower honey.

Table 1. Honey yield (HY) per colony and sugar consumption (SC) of groups per year
Çizelge 1. Grupların koloni başına bal verimi (HY) ve şeker tüketimleri (SC)

	Groups				
	I	II	III	IV	V
HY (kg)	11.4±2.30 ^{Aa}	14.2±1.25 ^{Aa}	16.4±1.47 ^{ABb}	20.0±1.78 ^{ABb}	21.4±1.38 ^{Bb}
SC (kg)	7.1±1.40	6.5±1.09	7.9±1.51	7.2±1.13	6.5±1.10

A, B; $P < 0.01$, a, b; $P < 0.05$

While groups I, II and III harvest honey once or twice per year, more frequent harvests (three or four times per year) were conducted by beekeepers in groups IV and V. The beekeepers in all groups stated that they produced both flower and honeydew honey. In addition to honey production, 15–20% of beekeepers in groups II and III groups, and 45% of beekeepers in groups IV and V produced pollen as well.

Beekeepers generally market their honey wholesale within a tin (about 27 kg of honey) to merchants or mediators at a price of about 90 US\$ per tin. Only 10% of the beekeepers market their honey in the jar. Beekeepers propose that the main problems with beekeeping in the region include the varroa mite, accommodation problems, the low price of honey, the cost of

moving colonies, diseases, inadequacy of government promotion and subvention.

In order to increase honey production, beekeepers in groups IV and V try to use different genotypes (Caucasian and Italian). Recently, some beekeepers tried using Carniolan honeybees in production and others were willing to try this honeybee, too. This pattern shows that some beekeepers imitate the practices of others, which is a typical traditionalist structure among beekeepers in many regions. During the production period, the beekeepers in Turkey start moving all over Turkey, from the Aegean and Mediterranean regions to the central and eastern Anatolia regions and then they come back to the Aegean region again. The beekeepers in groups IV and V observed that the performances of

Caucasian and Carniolan honeybees in central Anatolia, eastern Anatolia and the Thrace region exceeded those of the Aegean ecotype. This observation, explained by the beekeepers in this study, supports the idea that a genotype cannot be expected to provide satisfactory results in all regions (Uçak Koç and Karacaoğlu, 2013).

In recent years, one of the most important issues all over the world has been wintering losses in beekeeping. In this survey, we noted that wintering losses (between 6% and 29%) increased as the year advanced (Table 2). This result is in agreement with a study conducted in the Aegean region by (Tunca and Çimrin, 2012). However, for the Aegean region, reported a higher mortality rate than was found by our study (Uçak Koç, 2014).

The main reasons for the high mortality rates could include the varroa mite and diseases resulting from the intensive migration rate and the

closed architectures of the bee yards. Furthermore, wintering weak colonies, more sunny and warm days and high differences between the day and night temperatures in winter (which causes the winter cluster narrowing and expansion) and also early physiological aging because of the larger number of daily flights in winter may be responsible (Neukirch, 1982; Remolina and Hafez, 2007; Uçak Koç, 2014). It was found a strong correlation ($r = -71$, $r = -69$) between over wintering ability of colonies and the varroa mite infestation rate (Akyol and Yeninar, 2011).

Wintering losses in this region may not be a real problem for beekeepers compared with beekeepers in the United States and other countries. However, in some periods of the year in which the pesticide usage is high in intensified agriculture, an increased mortality rate is typically observed.

Table 2. Winter colony losses of the groups (%)

Çizelge 2. Grupların koloni kış kayıpları (%)

Years	Groups					Overall
	I	II	III	IV	V	
2009–2010	7.2	10.0	8.7	6.0	7.5	8.2
2010–2011	11.4	11	8.6	7.0	8.7	9.3
2011–2012	29.4	12.4	12.8	9.8	11.4	13.8
Overall	16.0	11.1	10.1	7.6	9.3	

One of the most important problems that the beekeepers mentioned was the accommodation locale and the closed nature of the apiaries during accommodation. Especially in the honeydew flow time, overcrowding of migratory beekeepers in the region increases these problems and creates negative effects for the local beekeepers. Overcrowding in the region results in sharing nectar resources, feeding colonies frequently and increasing the likelihood of contamination from rapine, varroa mites and diseases. As a result, the productivity of colonies decreases. In Turkey, there are approximately 7 colonies per square kilometer. At certain times in the Aegean region, however, because of the high migration rate, the colony density increases. However, in Canada and Australia, where the most honey is produced per colony, the number of colonies per square

kilometer is far below one (Karacaoğlu and Uçak Koç, 2007).

4. Conclusions and Recommendations

In conclusion, the rich nectar sources (pine forests and flowers) and mild climatic conditions made the Aegean region a center of attraction for beekeepers. The migration of a significant number of beekeepers to certain places in the region irregularly results in important problems such as accommodation, crowded bee yards, varroa mites and diseases and also results in shortages of the nectar resources. In addition, intensive pesticide usage in industrial agriculture in the lowlands and weak colony wintering because of mild winters in the region sometimes results in high colony losses in the region. In order to address the problems plaguing beekeepers in the region, reorganizing

the baiting of beekeepers, providing support only to the beekeepers make a living from beekeeping (groups IV and V) and creating an organizational structure among beekeepers are suggested.

References

- Akyol E ve Yeninar H (2011). The effects of varroa (*Varroa destructor*) infestation level on wintering ability and survival rates of honeybee (*L.*) colonies. *Journal of the Faculty Veterinary Medicine, Kafkas University*, 17 (3): 507-509.
- Anonymous (2016). www.fao.org. Erişim Tarihi: 05.09.2016.
- Caron DM and Hubner J (2001). Winter loss survey. *American Bee Journal*, 141:778-779.
- Danka RG and Villa JD (2000). A survey of tracheal mite resistance levels in U.S. commercial queen breeder colonies. *American Bee Journal*, 140 (5):405-407.
- Doğaroğlu M, Özder M ve Polat C (1992). To compare their performance important honeybee race and ecotype of Turkey (*Apis mellifera L.*) at the thrace conditions. *Doğa Tr. J. of Veterinary and Animal Sciences*, 16:403-414.
- Gençer HV ve Karacaoğlu M (2003). The brood rearing activities and honey productions of caucasian race (*Apis mellifera caucasica*) and reciprocal crosses of Caucasian and Aegean ecotype of Anatolian honey bee (*Apis mellifera anatoliaca*) in Aegean conditions. *Yüzüncü Yıl University, Journal of Agricultural Science (J. Agric. Sci.)*, 13 (1):61-65.
- Güler A ve Kaftanoğlu O (1999). Morphological characters of some important races and ecotypes of Turkish honeybees (*Apis mellifera L.*)-II. *Tr. J. of Veterinary and Animal Sciences*, 23(3):577-581.
- Karacaoğlu M (2005). Morphological characters of Aegean ecotype of Anatolian bee (*A. m. anatoliaca*) and Italian race (*A. m. ligustica*) X Aegean ecotype crossbreed. *Adnan Menderes University, Journal of Agricultural Faculty*, 1 (2):41-46.
- Karacaoğlu M ve Uçak Koç A (2007). Ege Bölgesi Arıcılığında Fırsatlar ve Kısıtlar. *Ege Bölgesi Arıcılık Semineri*, 15-16 Şubat, s:25-32. İzmir.
- Lawal OA and Banjo AD (2010). Appraising the beekeepers knowledge and perception of pests problem in beekeeping business at different ecological zones in south western Nigeria. *World Journal of Zoology*, 5(2):137-142.
- Uçak Koç A ve Karacaoğlu M (2010). Bozdoğan'ın Arıcılık Potansiyeli. *Bozdoğan Çevre ve Kültür Sempozyumu*, 13-15 Mayıs cilt 2, s: 135-149. Aydın.
- Uçak Koç A ve Karacaoğlu M (2013). Colony development of Caucasian (*A. m. caucasica*), Italian (*A. m. ligustica*) and Aegean ecotype of Anatolian (*A. m. anatoliaca*) honeybee races and their crosses under Aegean region conditions. *e-TRALLEIS*, 2013, 1: 28-35.
- Neukirch A (1982). Dependence of the lifespan of the honeybee (*Apis mellifera*) upon flight performance and energy consumption. *J Comp Physiol B.*, 146: 35-40.
- Tunca Rİ ve Çimrin T (2012). Kırşehir ilinde bal arısı yetiştiricilik aktiviteleri üzerine anket çalışması. *Iğdır Üniversitesi Fen Bilimleri Enstitüsü Dergisi*, 2(2):99-108.
- Remolina SC, Hafez DM, Robinson GE and Hughes KA (2007). Senescence in the worker honey bee *Apis mellifera*. *J. Insect Physiol* 53:1027-1033.
- Uçak Koç A (2014). Effects of altitude and beehive bottom board type on wintering losses of honeybee colonies under subtropical climatic conditions. *Spanish Journal of Agricultural Research*, 12(1): 151-158.