



## Determination Of Antifungal Activity and Phenolic Compounds Of Endemic *Muscari aucheri* (Boiss.) Baker Extract

Abdurrahman ONARAN<sup>1\*</sup> Mustafa BAYRAM<sup>2</sup>

<sup>1</sup>Department of Plant Protection, Faculty of Agriculture, Gaziosmanpaşa University, Tokat

<sup>2</sup>Department of Food Engineering, Faculty of Engineering and Natural Sciences, Gaziosmanpaşa University, Tokat

\*e-mail: abdonaran@hotmail.com

Alındığı tarih (Received): 20.06.2017

Kabul tarihi (Accepted): 22.02.2018

Online Baskı tarihi (Printed Online): 04.04.2018

Yazılı baskı tarihi (Printed): 30.04.2018

**Abstract:** Antifungal activity and phenolic compounds of the methanol extract (flower + peduncle) derived from *Muscari aucheri* (Boiss.) Baker that grows endemically in Turkey were identified against five different plant pathogens [*Fusarium oxysporum f. sp. cucumerinum*, *Alternaria solani*, *Verticillium dahliae*, *Rhizoctonia solani*, *Botrytis cinerea*]. Methanol extract of *M. aucheri* was obtained from the aerial parts (flower + peduncle). Antifungal activity studies were performed by examining the obtained extract against plant pathogens at the final concentrations of 2.5, 5, 10 and 20 mg/ml doses using the agar plate method. Mycelial growth inhibition and LD<sub>50</sub> doses of the extract to the pathogens were determined. Significant levels of antifungal activities were observed at all the doses used in the extracts. At the same time, 100% inhibition were observed at 10 and 20 mg/mL doses used against the pathogens. Individual phenolic compound rutin (693,2 ug/g) was found in the extract at its highest level. Total phenolic compound and monomeric anthocyanin amounts were determined. Results from these findings suggest that phenolic compounds in natural antifungal agents may offer positive results in the control of plant pathogens.

**Keywords:** *Muscari aucheri*, phenolic compounds, antifungal activity, plant pathogens

### Endemik *Muscari aucheri* (Boiss.) Baker. Ekstraktının Antifungal Aktivitesi ve Fenolik Bileşiklerinin Belirlenmesi

**Öz:** Türkiye’de endemik olarak yetişen *Muscari aucheri* (Boiss.) Baker bitkisinden elde edilen metanol ekstraktının (Çiçek + çiçek sapı) beş farklı bitki patojenine [*Fusarium oxysporum f. sp. cucumerinum*, *Alternaria solani*, *Verticillium dahliae*, *Rhizoctonia solani*, *Botrytis cinerea*] karşı antifungal aktivitesi ve bazı bireysel fenolik bileşikler belirlenmiştir. *M. aucheri*’nin metanol ekstraktı toprak üstü aksamından elde edilmiştir. Elde edilen ekstraktın son konsantrasyonu 2.5, 5, 10 ve 20 mg/mL dozlarında agar petri yöntemi kullanılarak bitki patojenlerine karşı antifungal aktivite çalışmaları yürütülmüştür. Ekstraktın patojenlere olan miselyum gelişim engellemeleri ve LD<sub>50</sub> dozları belirlenmiştir. Kullanılan bütün dozlar ve ekstraktta önemli düzeyde antifungal etkiler gözlenmiştir. Patojenlere karşı kullanılan 10 ve 20 mg/mL dozunda %100 engellemeler gözlenmiştir. Ekstraktta en yüksek bireysel fenolik bileşik rutin (693,2 µg/g) olarak bulunmuştur. Toplam fenolik bileşik ve monomerik antosiyanin miktarları belirlenmiştir. Bu sonuçlara göre, doğal antifungal maddelerin içerdiği fenolik bileşiklerin bitki patojenlerinin kontrolünde ümitvar sonuçlar ortaya çıkaracağı belirlenmiştir.

**Keywords:** *Muscari aucheri*, fenolik bileşikler, antifungal aktivite, bitki patojenleri

#### 1. Introduction

All plants produce many numbers of phenolic components in their metabolisms as seconder metabolites. Therefore, plants contain a wide range of phenolic in different quality and amount (Rauha et al., 2000; Proestos et al., 2006; Xia et al., 2011).

The usage of plants as bio-pesticides due to the rich antifungal metabolites they contain dates back to the start of human history. Antifungal effects of plants and plant based products become clearly evident every passing day. Antifungal metabolites obtained from plants present unique benefits with generally no side effects. Nowadays, it has emerged that side effects of pesticides used

against plant pathogens are harmful to the environment and people. Therefore, researchers for alternative management methods that will reduce the usage of pesticides to minimum come in to prominence. Following the researches, compounds and essential oils that plants contain are discovered to show antifungal (Gökce et al., 2006; Kordali et al., 2013; Bucchini et al., 2015; Elansary et al., 2016), antibacterial (Deans and Ritchie 1987; Hammer et al., 1999; Xia et al., 2011), herbicidal (Kordali et al., 2009; Verdeguer et al., 2011; Yılar et al., 2013, nematicidal (Elbadri et al., 2008; Kepenekci and Saglam 2015) and antiviral (Abad et al., 1999; Vijayan et al., 2004) activities.

*Muscari aucheri* (Boiss.) Baker. is a perennial bulbous plant is known as Grape Hyacinth. It is an endemic plant originally from Turkey (Uranbey, 2010). *M. aucheri* is usually less than 10 centimetres tall. There are two or three leaves per bulb, peduncle and a flower. The flowers are arranged in a raceme (Mathew, 1987). *Fusarium oxysporum* (Schlechtend.:Fr.) f. sp. *cucumerinum* (Owen) Snyder & Hansen [FOC], *Alternaria solani* Sorauer, *Verticillium dahliae* Kleb., *Rhizoctonia solani* J.G. Kühn and *Botrytis cinerea* Pers., cause damages in cucumber, tomato, potato and strawberry plants in the world respectively. FOC is the causal agent for fusarium wilt disease in cucumbers in many parts of the world. It causes serious economic damages in cucumbers. This pathogen is particular to cucumber plant (Jenkins and Wehner., 1983). The factor of the disease called early blight that can widely be found on tomato plants is *A. solani* (Agrios, 1988). *V. dahliae* is a soil born pathogen that causes *verticillium* wilt disease in tomato plants (Diwan et al., 1999). *R. solani* is the effects of the disease called soft rotteness in potatoes roots and tubers and causes serious crop losses in potatoes (Carling et al 1989). *B. cinerea* is the effect that causes grey mold disease in strawberries around the world where strawberry production is made (Grabke et al 2014).

This study aims that to show antifungal activities of individual phenolic compounds obtained from methanol extract of *M. aucheri*

plant that grows endemically in Turkey and to identify their effects to control diseases caused by FOC, *A. solani*, *V. dahliae*, *R. solani* and *B. cinerea* pathogens.

## 2. Methodology

### 2.1. Plant Material

*Muscari aucheri* plants were collected from Tokat, Turkey in April, 2016. Aerial parts of plant (flower and peduncle) were used for the experiment. Plants were washed by sterile water and then dried in room temperatures under shadow. Dried plants materials were passed through a mill to separate small pieces.

### 2.2. Fungi Cultures

Plant pathogens (Table 1) were obtained from stock cultures at Department of Plant Protection Gaziosmanpaşa University, Fungi cultures were developed at 20 mL potato dextrose agar (PDA) on petri dish (90 mm) and kept at 22±2 °C during 7 days and these fungi were used for the experiment. The abbreviations used in the article of plant pathogens are given in Table 1.

### 2.3. Plant Extracts

The aerial parts of *M. aucheri* (100 g) were weighted and put 1 lt jar. The methanol (Merck, Germany, 1.06007.2500) was added to cover the plant materials. These were incubated on an orbital shaker (Lab. Corporation Group, Model SI-300) (120 rpm) at 30°C for 3 days. The methanol was evaporated to dryness in the rotary evaporator (Heildolph Group, Model Hei-Vap-Precision) at 40°C. These dried plant materials were reconstituted solution by 5% dimethyl sulfoxide (DMSO) (Onaran and Yılar 2012).

### 2.4. *In vitro* Antifungal Activity

The antifungal activities were determined by using agar plate methods (Nwosu and Okafor, 1995). PDA [95mL (w/v)] was autoclaved and kept at 40°C. Dried plant extracts were weighted 250, 500, 1000 and 2000 mg and reconstituted solution by 5% dimethyl sulfoxide (DMSO), and then added to PDA media (95 mL). The final

concentrations were arranged to 2.5, 5, 10 and 20 mg/mL and transferred to PDA petri plate (60 mm) (~10mL plate<sup>-1</sup>). The mycelium disc (5 mm in diameter) from 7-day-olds fungi cultures were transferred to petri plates. Then incubated at 22±2 °C (Memmert Group, Model-High Precision

Incubator) during 7 days and were recorded growth of fungi daily (Onaran and Yılar, 2012). The recommended dose of commercial fungicide (80% Thiram) was used as a positive control. DMSO (%5) was used as a negative control. The experiment was set up 3 replicate and twice.

**Table 1.** The list of plant pathogens, their abbreviations and the name of plants which pathogens were isolated

**Çizelge 1.** Patojenlerin izole edildiği bitkilerin isimleri ve kısaltmaları, bitki patojenlerinin listesi

Plant Pathogens	Abbreviations	Isolated
<i>Fusarium oxysporum f. sp. cucumerinum</i> J.H. Owen	Foc	Cucumber
<i>Alternaria solani</i> (Ell. And G. Martin)	As	Tomatoes
<i>Verticillium dahliae</i> (Kleb.)	Vd	Tomatoes
<i>Rhizoctonia solani</i> Kühn	Rs	Potatoes
<i>Botrytis cinerea pers.:Fr</i>	Bc	Strawberry

The percentage of mycelial growth inhibition was calculated accordingly the formula mentioned by Pandey et al. (1982):

$$I=100 \times (dc-dt)/dc$$

I: Mycelial growth inhibition

dc: the mycelial growth in control

dt: the mycelial growth in treatment

## 2.5. Determination of Some Individual Phenolic Compounds

The gallic acid, 4 hydroxybenzoic acid, vanillic acid, cafeic acid, ferulic acid, rutin, cistic acid, rozmarinic acid, quercetin, gentisic acid were purchased from Sigma-Aldrich Chemical Co. (United States). The analysis was carried out by LC 20AT pump (Shimadzu), SPD-M20A model DAD detector (280 nm) and HPLC system with CTO-20AC model column oven; for separation, Dionex, (150x4.60 mm, 3 µm) C16 reversed phase filler column were used. The solvent A (2,5% methanol with formic acid), solvent B (2,5% deionized water with formic acid) and solvent C (acetonitrile) were used for mobile phase at HPLC. 10 mL extract samples which used for analysis were filtered by 0,45 µm membrane filter. The flow rate was 1 mL/min, injection volume was 20 µl and the column temperature was set at 30 °C. The gradient elution program was given in Table 2.

## 2.6. Determination of Total Phenolic Compounds

Total phenolic compounds were determined by using Folin-Ciocalteu method (Singleton and

Rossi, 1965). Results were expressed as mg gallic acid equivalents (GAE) per litre of sample extracts (µg GAE/g).

**Table 2.** The gradient elution program for determination of phenolic compounds

**Çizelge 2.** Gradient elüsyon programı ile fenolik bileşiklerin belirlenmesi

Time (min.)	Solvent A (%)	Solvent B (%)	Solvent C (%)
0	5	95	0
9	10	90	0
27	15	85	0
29	0	85	15
40	0	70	30
50	0	45	55
55	0	0	100

## 2.7. Determination of Total Monomeric Anthocyanin

Total anthocyanin content of extracts were evaluated by pH-differential method described by Giusti and Wrolstad (2001). By this method, buffer of 0.025 M KCl (pH 1.0) and buffer of 0.4 M CH<sub>3</sub>COONa (pH 4.5) dilutions were equilibrated for 15 min. The absorbance of each solution was measured at 520 nm and 700 nm.

Results were calculated by the following formula and expressed as µg malvidin 3-glycoside equivalent per gram.

$$A = (A_{\lambda_{520}} - A_{\lambda_{700}})_{\text{pH 1.0}} - (A_{\lambda_{520}} - A_{\lambda_{700}})_{\text{pH 4.5}}$$

$$\text{TA (mg/kg)} = A \times \text{MA} \times \text{SF} \times 1000 / \epsilon \times 1$$

A: absorbance, Malvidin-3-O-glycoside

molecular weight (MA): 493.5 g/mol/L;

The dilution factor (DF);

$\epsilon$ , molar absorptivity (28.000)

## 2.8. Statistical Analysis

Data were analysed by using One Way procedure of ANOVA (Windows version of SPSS, release 15.00). Differences among concentrations were compared with using DUNCAN Multiple Range Test of  $p < 0.05$ . The probit analysis of the data derived in consequence of the tests was performed through POLO Plus 1.0 computer program and the values of LD<sub>50</sub> were calculated.

## 3. Results and Discussion

### 3.1. Some Individual Phenolic Compounds, Total Phenolic Compounds, Total Monomeric Anthocyanin Content

HPLC chromatogram of the standart (280 nm) and the extract (280 nm and 330 nm) were shown in figure 1. The amount of individual phenolic compounds were shown Table 3. The highest individual phenolic compound was found to rutin (693,2 µg/g) from *Muscari aucheri*. This was followed by ferulic acid (377,9 µg/g), rosmarinic acid (127,9 µg/g) respectively. Phenolic compounds from the aerial part (flower + peduncle) of *M. aucheri* were determined first time with this study. Gallic acid, vanillic acid and cistic acid were not determined from *M. aucheri* plant extract (Table 3). In a study on the antioxidant capacity of this plant extract was showed that the amount of antioxidant capacity was found to be 3,164 (bulb) to 1,145 (leaf) mmol Trolox Equiv/L (Yıldırım 2013).

The total phenolic compound content was determined 1882,13 µg GAE/g from *M. aucheri* extract. In addition, the amount of total monomeric anthocyanin was calculated as 52,34

µg/g malvidin-3-O-glycoside equivalent. Phenolic compounds in plants consist of phenolic acids, flavonoids with small molecules and mostly essential compounds. Antifungal and antibacterial (Aziz et al., 1998; Rauha et al., 2000) and antiviral (Cline et al; 1969; Özcelik et al., 2011) activities of phenolic compounds of plants were reported from many studies. Phenolic acid such as rosmarinic acid and polyphenols have synergistic effects (Choi et al., 2002). The extracts of *M. aucheri* have high level of phenolic acids had antifungal activity.

**Table 3.** The amount of individual phenolic compound from extract

**Çizelge 3.** Ekstratın bireysel fenolik madde miktarı

Retention Time	R <sup>2</sup> value	Phenolic Compounds	Amount (µg/g)
5.607	0,999	Gallic Acid	nd*
12.068	0,998	4hydroxybenzoic Acid	83,0
13.719	0,995	Gentisic Acid	99,4
14.901	0,999	Vanillic Acid	nd
17.432	0,997	Cafeic Acid	39,8
26.902	0,996	Ferulic Acid	377,9
34.952	0,998	Rutin	693,2
37.119	0,999	Cistic Acid	nd
38.575	0,999	Rosmarinic Acid	127,9
44.980	0,999	Quercetin	8,8

\*nd: not detected


### 3.2. In vitro Antifungal Activity

Antifungal activities were found differently rate for each plant pathogens. All doses of plant extract were showed antifungal activity noticeably. This activity has also increased with the rise of the amount of doses. The highest dose (20 mg/mL) of plant extract was observed 100% mycelium inhibition rate on *Alternaria solani*, *Verticillium dahliae*, *Rhizoctonia solani* and *Botrytis cinerea*. This was followed by 88% of inhibition on *Fusarium oxysporum f. sp. cucumerinum* (Table 4). These results were compared with positive control, similar inhibition rates were found. In the study, LD<sub>50</sub> values of plant extract was calculated against plant pathogens (Figure 2). According the value of the LD<sub>50</sub>, *V. dahliae* was found most susceptible pathogen against plant extract. This was followed

by *A. solani*, *R. solani*, *B. cinerea* ve *Fusarium oxysporum f. sp. cucumerinum*. All doses of plant extract compare with negative control was showed antifungal effect on plant pathogens (Figure 3). Mycelium growth was found to stop completely dependent on the doses increases. Even though lowest dose was showed limited mycelium growth, it was stopped after a certain period time. According the difference of fungi species, there have observed that differential such as aerial mycelium growth and inhibition of

sporulation formation (Figure 3). One of the similar study reported by Yıldıırım (2013), the leaf and bulb of *M. aucheri* was collected from Munzur valley of Tunceli.

The leaves and bulbs were used for obtaining methanol, n- hexane, ethanol and water extracts against identifying antifungal effect. The dose of 2.5% (SDA media) of extracts were used against *Coriolus versicolor* and found low antifungal activities.



**Figure 1.** HPLC chromatogram of the phenolic standards at 280 nm (A); HPLC chromatogram of the extract at 280 and 330 nm (B); Gallic acid (1), 4 hydroxybenzoic acid (2), Gentisic acid (3), Vanillic acid (4), Caffeic Acid (5), Ferulic Acid (6), Rutin (7), Cistic acid (8), Rozmarinic acid (9), Quercetin (10)  
**Şekil 1.** Fenolik standartların 280 nm’de HPLC kromatogramı (A); Ekstratın 280 ve 330 nm’de HPLC kromatogramı (B); Gallik asit (1), 4 hidroksibenzik asit (2), Gentisik asit (3), Vanillik asit (4), Kafeik asit (5), Ferulik asit (6), Rutin (7), Sisik asit (8), Rozmarinic asit (9), Kuersetin (10)

Methanol and n-hexane extracts had not showed any antifungal activities but the leaf of ethanol extracts was found mycelium inhibition at 44.40% the leaf of water extract, 28.80% the leaf of water extract, 10.40 % the bulb of ethanol extract and 12,80 % the bulb water extracts

respectively. According to this finding, different parts of extracts of *M. aucheri* against different plant pathogens have shown antifungal activities. This difference may be due to the fact that the same plant species was grown in different areas.

In similar studies, the test organisms which used our study have been tested different plant extracts for identifying antifungal activities by several researchers such as *Fusarium oxysporium* (De Rodriguez et al., 2005; Çakır et al., 2005; Onaran and Yılar, 2012), *Alternaria solani* (Çakır et al., 2005; Yanar et al., 2011) *Verticillium dahliae* (Arslan ve Derviş 2010; Onaran and Yılar, 2012) *Rhizoctonia solani* (Alkhail, 2005, De Rodriguez et al., 2005, Onaran and Yılar 2012) *Botrytis cinerea* (Alkhail, 2005; Soylu et al., 2010). In this study, the methanol extracts of flower and peduncle parts of endemic *M. aucheri* has been studied first time identifying antifungal activities


according to literatures. This aspect increases the originality of the study.

**Table 4.** Mycelial growth inhibition of fungi at different doses of *Muscari aucheri* (%)

**Çizelge 4.** *Muscari aucheri*'nin farklı dozlarında fungusların miselyum gelişim engellemeleri (%)


Doses (mg/mL)	Foc	As	Vd	Rs	Bc
C-	0	0	0	0	0
2,5	20	48	50	28	22
5	56	66	75	52	58
10	85	95	100	88	84
20	88	100	100	100	100
C+	100	100	100	100	100

C-: Negative control; C+: Positive control



**Figure 2.** LD<sub>50</sub> values of plant extract against used plant pathogens

**Şekil 2.** Kullanılan bitki patojenlerine karşı bitki ekstraktlarının LD<sub>50</sub> değerleri



**Figure 3.** The growth of mycelium of pathogens against the extract of *Muscari aucheri* (The photo was taken at the end of 7<sup>th</sup> day).

**Şekil 3.** *Muscari aucheri* ekstratının patojenlerin miselyum gelişimi üzerine etkisi (Fotoğraf 7. Günün sonunda çekilmiştir.)

#### 4. Conclusions

Plants have antifungal compounds such as flavonoids and phenolic compounds therefore plants have always been of interest the researchers. Each day, new compounds find from plants are identified and used in scientific studies. As a result, new identified compounds of plant structures may replace the synthetic chemicals which heavily used in industry, agricultural practice and food industry. Thus, this allows to find natural substances that are less harmful or harmless to human, environment and non-target organisms. With this study, the flower and peduncle of endemic *Muscari aucheri* plant in Turkey were done to collected and identified of phenolic compounds and antifungal activities. According to result, this compounds and extracts give promising result to control test pathogens so they can be replaced the synthetic chemicals that are used to control them.

#### References

- Abad MJ, Bermejo P, Palomino SS, Carrasco L, Chiriboga X (1999). Antiviral activity of some South American medicinal plants. *Phytotherapy Research*, 13(2):142-146.
- Agrios GN (1988). *Plant pathology*, Academic Press Limited 24-28 oval, London NW1, 7DX, p. 803.
- Alkhail AA (2005). Antifungal activity of some extracts against some plant pathogenic fungi. *Pakistan Journal of Biological Sciences*. 8(3):413-417.
- Arslan M, Dervis S (2010). Antifungal activity of essential oils against three vegetative compatibility groups of *Verticillium dahliae*. *World Journal of Microbiology and Biotechnology*. 26:1813-1821.
- Aziz NH, Farag SE, Mousa LA, Abo-Zaid MA (1998). Comparative antibacterial and antifungal effects of some phenolic compounds. *Microbios*. 93:43-54.
- Bucchini A, Ricci D, Messina F, Marcotullio MC, Curini M, Giamperi L (2015). Antioxidant and antifungal activity of different extracts obtained from aerial parts of *Inula crithmoides* L. *Natural Product Research*. 29:1173-1176.
- Carling DE, Leiner RH, Westphale PC (1989). Symptoms, signs and yield reduction associated with *Rhizoctonia* disease of potato induced by tuber borne inoculum of *Rhizoctonia solani* AG-3. *The American Journal of Potato Research*. 6:639-697.
- Choi CW, Kim SC, Hwang SS, Choi BK, Ahn HJ, Lee MY, Park SH, Kim SK (2002). Antioxidant activity and free radical scavenging capacity between Korean medicinal plants and flavonoids by assay-guided comparison. *Plant Science*. 163:1161-1168.
- Cline JC, Nelson JD, Gerzon K, Williams RH, DeLong DC (1969). In Vitro Antiviral Activity of Mycophenolic Acid and Its Reversal by Guanine-Type Compounds. *Applied Microbiology*. 18: 14-20, 1969.
- Çakir A, Kordali S, Kilic H, Kaya E (2005). Antifungal properties of essential oil and crude extracts of *Hypericum linarioides* Bosse. *Biochemical Systematics and Ecology*. 33:245-256.
- De Rodriguez DJ, Hernandez-Castillo D, Rodriguez-Garcia R, Angulo-Sanchez JL (2005). Antifungal activity in vitro of *Aloe vera* pulp and liquid fraction against plant pathogenic fungi. *Industrial Crops and Products*. 21:81-87.
- Deans SG, Ritchie G (1987). Antibacterial properties of plant essential oils. *International Journal of Food Microbiology* 5, 165 – 180.
- Diwan N, Fluhr R, Eshed Y, Zamir D, Tanksley SD (1999). Mapping of Ve in tomato: a gene conferring resistance to the broad-spectrum pathogen, *Verticillium dahliae* race 1. *Theoretical and Applied Genetics*. 98(2):315-319.
- Elansary HO, Yessoufou K (2016). *In vitro* antioxidant, antifungal, and antibacterial activities of five international *Calibrachoa* cultivars. *Natural Product Research*. 30:1339-42.
- Elbadri GAA, Lee DW, Park JC, Yu HB, Choo HY, Lee SM, Lim TH (2008). Nematocidal Screening of essential oils and herbal extracts against *Bursaphelenchus xylophilus*. *Plant Pathology Journal*. 24(2): 178-182.
- Giusti MM, Wrolstad RE (2001). Characterization and measurement of anthocyanins by UV-visible spectroscopy. *Current protocols in food analytical chemistry*. New York: John & Wiley, Inc. P F1.2.1-F1.2.13.
- Gökçe A, Whalon ME, Çam H, Yanar Y, Demiştaş I, Gören N (2006). Plant extract contact toxicities to various developmental stages of Colorado potato beetles (Coleoptera: Chrysomelidae). *Annals of Applied Biology*. 149:197-202.
- Grabke A, Fernández-Ortuño D, Amiri A, Li X, Peres N. A, Smith P, Schnabel G (2014). Characterization of iprodione resistance in *Botrytis cinerea* from strawberry and blackberry. *Phytopathology*. 104:396-402.
- Hammer KA, Carson CF, Riley TV (1999). Antimicrobial activity of essential oils and other plant extracts. *Journal of Applied Microbiology*. 86:985 – 990.
- Jenkins JR, Wehner TC (1983). Occurrence of *Fusarium oxysporum* f. sp. *cucumerinum* on greenhouse-grown *Cucumis sativus* seed stocks in North Carolina. *Plant Disease*. 67:1024-1025.
- Soylu EM, Kurt Ş, Soylu, S (2010). In vitro and in vivo antifungal activities of the essential oils of various plants against tomato grey mould disease agent *Botrytis cinerea*. *International Journal of Food Microbiology*. 143:183-189.
- Kepenekçi I, Sağlam HD (2015). Extracts of some indigenous plants affecting hatching and mortality in the root-knot nematode [*Meloidogyne javanica* (Treub) Chitwood]. *Egyptian Journal of Biological Pest Control*. 25(1): 39-44.
- Kordali Ş, Çakır A, Akcin TA, Mete E, Akcin A, Aydın T, Kılıç H, (2009). Antifungal and herbicidal properties of essential oils and n-hexane extracts of *Achillea*

- gypsicola* Hub-Mor. and *Achillea biebersteinii* Afan. (Asteraceae). Industrial Crops and Products. 29:562-570.
- Kordali Ş, Usanmaz A, Cakir A, Cavusoglu A, Ercisli S (2013). In Vitro antifungal effect of essential oils from *Nepeta meyeri* Benth. Egyptian Journal of Biological Pest Control. 23(2):209-213.
- Mathew B (1987). The Smaller Bulbs, London: B.T. Batsford, ISBN 978-0-7134-4922-8, p. 126.
- Nwosu MO, Okafor JI (1995). Preliminary studies of the antifungal activities of some medicinal plants against *Basidiobolus* and some other pathogenic fungi. Mycoses 38:191-195.
- Onaran A, Yılar M (2012). Antifungal activity of *Trachystemon orientalis* L. aqueous extracts against plant pathogens. Journal of Food, Agriculture and Environment. 10:287-291.
- Özcelik B, Kartal M, Orhan I (2011). Cytotoxicity, antiviral and antimicrobial activities of alkaloids, flavonoids, and phenolic acids. Pharmaceutical Biology. 49, 396-402.
- Pandey DK, Tripathi NN, Tripathi, RD, Dixit SN, (1982). Fungitoxic and phytotoxic properties of essential oil of *Hyptis suaveolens*. Z. Pflanzenk, 89, 344-349.
- Proestos C, Boziaris IS, Nychas GJE, Komaitis M, (2006). Analysis of flavonoids and phenolic acids in Greek aromatic plants: investigation of their antioxidant capacity and antimicrobial activity. Food Chemistry. 95:664-671.
- Rauha JP, Remes S, Heinonen M, Hopia A, Kahkonen M, Kujala T (2000). Antimicrobial effects of Finnish plant extracts containing flavonoids and other phenolic compounds. International Journal of Food Microbiology. 56:3-12.
- Singleton VL, Rossi JA (1965). Calorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. American Journal of Enology and Viticulture. 16:144-158.
- Uranbey S (2010). Stimulating effect of different basal media and cytokinin types on regeneration of endemic and endangered *Muscari aucheri*. Archives of Biological Sciences. 62(3):663-667.
- Verdeguer M, Blazquez MA, Boira H (2011). Chemical composition and herbicidal activity of the essential oil from a *Cistus ladanifer* L. population from Spain. Natural Product Research. 26:1602-1609.
- Vijayan P, Raghu C, Ashok G, Dhanaraj SA, Suresh B (2004). Antiviral activity of medicinal plants of Nilgiris. Indian Journal of Medical Research. 120:24-29.
- Williams AH (1962). Enzyme inhibition by phenolic compounds. In Pridham JB (ed) Enzyme chemistry of phenolic compounds. Pergamon Press, New York. Pp:87-95.
- Xia DZ, Yu XF, Zhu ZY, Zou ZD (2011). Antioxidant and antibacterial activity of six edible wild plants (*Sonchus spp.*) in China. Natural Product Research. 25: 1893-1901.
- Yanar Y, Gokce A, Kadioglu I, Cam H, Whalon M (2011). In vitro antifungal evaluation of various plant extracts against early blight disease (*Alternaria solani*) of potato. African Journal of Biotechnology. 10:8291-8295.
- Yılar M, Bayan Y, Aksit H, Onaran A, Kadioglu I, Yanar Y (2013). Bioherbicidal effects of essential oils isolated from *Thymus fallax* F., *Mentha dumetorum* Schult and *Origanum vulgare* L. Asian Journal of Chemistry. 25:4807-4811.
- Yıldırım NC, Paksoy MY, Yüce E, Yıldırım N (2013). Total Antioxidant status and Antifungal Activities of Endemic Geophytic Plants Collected from Munzur Valley in Tunceli, Turkey. Digest Journal of Nanomaterials and Biostructures. Vol. 8 (1): 403-408.